

31

Dysentery

17

Thomas Old

admitted March 14th 1821.

1794

12th

12th

1794

last the
telling the
a full month
by and carefully
that in the
the present the
from which
last of January
house, that was
which is still
by the day
which has been
of the last
by way of the
a perfect
to young men
at home
which was
let in the
which is the

A Dissertation on Dysentery.

1

Amidst the many subjects offered the candidate, upon which to display his abilities or explain his views; there is not one on which we feel ourselves capable either of advancing new ideas, or satisfactorily and usefully discussing those which now obtain the ascendancy. Shall we succeed in attaining useful andivociry, we shall be better pleased than by the collision of opposition, to elicit the applause of some, whilst we deservedly receive the contempt of others for our heedless temerity. We could perhaps advance wild unsupported theories, but no ingenuity of defence ought to save them from merited disapprobation. In medicine we want truth not sophistry tho' dropped never so bewitchingly, and every honest man will contribute his best endeavours for the support of the former and the abolition of the latter in all our disputations and contests, whatever they may be. To propose theories, or to pretend to originality, would argue a profundity of research, which tho' we would aspire to, to fancy ourselves possessed of, would be presumption in the highest degree unpardonable.

Without any farther prefatory remarks, we will now observe that we have chosen Dysentery as the subject of our essay. In this selection, we have been influenced by the frequent occurrence of the disease

and its occasion
much new, or y
shall we succeed
been a correct
think best adapt
fortunate. Show
the difficulties
labours under is
for the many
following pages
of dysentery, top
employed in its
an evidence of the
opinions are und
wish them to hold
willed to. We an
to have consulted
disorder, we might
practice and perh
was precluded and

and its occasional great mortality, rather than with the hope of saying much new, or greatly adding to our stock of knowledge on this subject. Shall we succeed in giving a clear and succinct view of what we deem a correct theory of this disease, and of that treatment which we think best adapted to its cure, we shall esteem ourselves peculiarly fortunate. Should we fail in our attempt, we trust that the insuperable difficulties which a young man, unaided by practical experience labours under in composition, will stand as a sufficient excuse for the many errors we may be guilty of. Our objects will be in the following pages so far as we are able, to give the causes and treatment of Dysentery, together with the modus operandi of the remedies employed in its removal, and their connexion with each other. We are conscious of the magnitude of our undertaking, but where our opinions are unsupported by sound and legitimate argument, we wish them to hold only that rank which all vague suppositions are entitled to. We are fully aware that had we time and opportunity to have consulted the many good works which we possess on this disease, we might have added much to the length of the following treatise and perhaps not a little to its merit; but from these we were precluded and we trust that shall we succeed in attaining

he objects we
Dysentery

"Pyre

late, relaps

this definition

say we have seen

syndromy has

until within a

constitutes the p

it was first des

only European ac

finism. He say

Remittent form

light and have

let to the contag

first American

There is to att

disease of a like

intensity, we sh

for the present p

the objects we have assumed, it will be all that is required of us.

Dysentery is defined by Cullen,

"Pyrexia contagiosa; dejectiones frequentes, mucosa vel sanguinea lenta, siccitas plerumque facilius alvinis; tormina; tenesmus."

This definition we find conveying a more correct idea of the disease than any we have seen, yet it is not unexceptionable. The contagiousness of Dysentery has been admitted on all sides from time immemorial until within a few years past, and even now with some few exceptions constitutes the prevailing doctrine on the Continent of Europe. By whom it was first denied we are unable certainly to say, tho' Dr. Hasley is the only European author with whom we have met, who advocates a contrary opinion. He says, that Dysentery neither in its simple, intermittent, or remittent form is contagious; that it is only so when combined with typhus and here he ascribes it not to the virus specific to Dysentery, but to the contagion of Fever. Dr. Chapman we believe to have been the first American physician who advanced a similar doctrine.

When we to attempt to explain why Dysentery with so many other diseases of a like character have been ranked under the head of contagion, we should be led into a disquisition infinitely too lengthy for the present paper. We will merely observe that it appears to us

to have arisen
between Contagion
been enough to
universal prom-
shly neglecting
facing the charac-
ter, but would
constitute Contagion
such decided as
agreed to convene
on property of property
that property would
property of more than
end of these classes
to medical science
kind to allow it; but
which is all that can
of Contagion
under particular laws
cases, in any situation

to have arisen from Physicians, not having sufficiently discriminated between Contagious, Atmospheric and Endemic diseases - it having hitherto been enough to establish the contagion of a disease, to ascertain its universal prevalence in particular neighbourhoods, districts or towns; wholly neglecting farther to investigate its remote cause. By this mode of fixing the character of a disease, not one belonging to either of the above classes, but would be found to possess all the qualities necessary to constitute contagion. We having ascertained that there really exist such diseases as Contagious, Atmospheric and Endemic, is all that is requisite to convince us, that it was very possible and probable from one property possessed in common by all, to range any disease having that property under the head of contagion. This equally shows us the necessity of more precisely ascertaining and defining the boundaries between each of these classes of diseases. A work on this plan would contribute much to Medical Science. We confess ourselves incompetent to the task, would our limits allow it, but we will venture to give a definition of each of them, which is all that our present purposes require.

1st Of Contagion. By a contagious disease we understand one which (under particular laws) may be propagated independent of all extraneous causes, in any situation and under any circumstances, it being only

surpassing
distance of
such is subject
deserving to a
catalogue.

2nd

reach all the
the atmosphere
or variety in
may occasion
the atmosphere
"the general" is
and experience
has produced
Pompanona by

The will not
have been the
diversity in these
by the double ge
any other cause.

necessary that the person to be infected be brought within a definite distance of the sick, still keeping it in view that this as every other general rule is subject to exceptions and anomalies; at the same time wholly denying to all diseases not possessed of these properties the character of contagion.

2nd Atmospheric Diseases. We would under this head rank all those diseases, which are produced by the insensible qualities of the atmosphere, denying the influence of its relative heat, moisture, density or rarity in their generation; and at the same time admitting that these may occasionally prove exciting causes. By the insensible qualities of the atmosphere we would mean the existence of some fluid, or mist, or "vis genies" in the air whose properties are not cognizable to the senses, and experience has shown that these may be various, as the diseases thus produced are numerous. Of this class our late winter disease, the *Varicella* is a prominent example.

We will now pass to our third class of Endemic Diseases, which have been the chief cause of our noticing the two preceding ones. Endemic diseases are those which attack many persons at the same time, and are excited by the insensible qualities of the atmosphere, as its heat, moisture, &c. &c. and any other cause of extension, prevalence, or to which many people are at the

situation of the French
 led differences of defini-
 tions. Thus we
 a meaning that was
 failed to it an obvious
 and gradually, &
 in defining and state-
 ments definition success-
 ful of this would be the
 fact and which we
 was in our opinion as
 set off by 7 with a
 and perfectly be-
 I generally have
 a deeply founded op-
 inion of matter, as the
 in contact with the
 and the different stag-
 es of the necessary

same time or in succession exposed, being no doubt also affected in their progress by the situation of the person placed under the influence of their remote cause; which differences of situation and other circumstances may prove the chief exciting causes. Thus much we have said, and now we will dismiss the subject only remarking that under this latter class we would place Dysentery, whose affinities to it are obvious and striking. We well know that Dysentery frequently occurs sporadically, but whenever it assumes its worst form, it is always as an Epidemic; and still tho' we cannot with propriety, according to the accepted definition annexed to the term *Enteritis*, class Dysentery under this head, yet this would be perfectly consonant to the meaning which we have afforded it, and which we flatter ourselves is altogether unobjectionable; a disease in our opinion arising from causes (to which all are subject tho' not all affected by) with the exception of our first and second classes, may with much propriety be thus ranked.

Dysentery has been divided by nosologists into many different species chiefly founded upon the preponderance of some one symptom, or the absence of another, as the appearance of Mucous instead of bloody stools, when combined with Intermittent Fever, &c. &c. Such divisions serve only to mark the different stages of the disease, or its mildness or severity, and being wholly unnecessary to practical forecision will be altogether

exhibited in the following
Symptoms. An
the feelings, and the other
often first appears.

Now and then
has been sometimes
and vomiting, which may
sometimes, nothing, the tem-
perature may be normal or
It sometimes comes on
which gradually increases
Pain is said to pro-
ceeding off the natural, and
and ultimately from ex-
The pain in this disease
is various, as plethoric and
rheumatic, pleuritic, &c.
and those.

The Pain in this d-
though it is to be regarded

neglected in the following sheets.

Symptoms. An attack of Dysentery is often preceded by alternate chills, flushings, and the other symptoms of Fever, the more generally the local affection first appears.

Now and then the primary symptoms are much more severe than have been mentioned; the Patient being affected with rigors, nausea and vomiting, which may even, (the more rarely) proceed to the ejection of stercoraceous matters, the termina and tendons being from the first inflamed, so they may be several days gradually progressing to that degree.

It sometimes comes on with a Diarrhea, the griping and frequency of the stools gradually increasing until it becomes a confirmed Dysentery.

Diarrhea is said to produce Dysentery by the repeated acid discharges washing off the natural mucus of the part, which thence becomes irritated and ultimately from erosion or lacerity ejects blood.

The pain in this disease is generally of the griping kind, tho' it may be various, as Coughorn and others describe it as occasionally resembling Stitches, Pleuritic pains, &c. &c. shooting into different parts of the abdomen and thorax.

The Fever in this disease is usually a Synocha and when it is a Typhus, it is to be regarded rather as a combination with Dysentery,

are produced by a
long mile, called the
early suburbs in a
very often the form is
be combined with the
the best affection and
both respect to the st
and finished with blue
and, and then again p
The birds sometimes
and being about the
Agave americana.
In Dyer's history the
by the alkali, it is in
old fields, an eva
a medium is always
When the lower part of
near the growing kind
the trunks and limbs are
colored line which is

or as produced by a protraction of the disease, than as an essential symptom. In very mild cases the fever is slight or wholly absent; here the disease usually subsides in a few days.

Very often the fever is of the Intermittent or Remittent type, or it may be combined with Intermittent Fever, and if the remission is distinct and the local affection moderate, we may conclude upon a manageable disease.

With respect to the stools, they are of various appearance, being sometimes only streaked with blood, or consisting of a little mucus mixed with blood, and then again pure blood being voided even from the commencement. The stools sometimes consist entirely of mucus, all appearance of blood being absent. This has been termed by authors the mucus stools or *Dysentery Alba*.

In Dysentery the natural forces are generally retained, and when they do appear, it is in the form of hard compact substances which are called scybala, an evacuation of which, whether produced by nature or medicine is always attended with a remission of all the symptoms.

When the lower part of the intestinal tube is affected, the pain is nearer the griping kind, particularly when situated in the rectum. The terminus and unctum are so excruciating as sometimes to produce a *Prolapsus Ani* which is always a troublesome symptom.

Therefore he is
silly head, the dis-
contented.

Proctorial was
advised to do
he had is more in
he sympathy existing
smaller entitled
his little confidence
him; but when con-
sidered, it may appear

The little sonnet
was first later, again
the patient affec-
he looks when atten-
of a good propo-

Others have been
not of the intention also
We suspected inward
The appearance of the

The nearer the stomach the pain is felt and the more it approaches the
 sickly kind, the disease is supposed to be situated higher up in the small
 intestines.

Practical writers have by their means and the appearance of the stools,
 endeavoured to discover the seat of the disease, placing it higher up when
 the blood is more intimately mixed with the faeces and vice versa; this from
 the sympathy existing between all the parts of the alimentary canal and the
 smaller intestines constantly changing their place, we are disposed to put
 but little confidence in the seat of the pain as indicating the seat of the
 disease; but when combined with the others which we think are founded in
 reason, it may assist us in forming an opinion.

The stools sometimes are of a dark, grumous appearance and have an
 offensive odor; again they have little or no smell; the former symptoms
 either portend approaching gangrene or show it actually to have taken place;
 the latter when attended with a corresponding mildness of the other symptoms
 afford a good prognosis.

Authors have been in the habit of mentioning the appearance of the internal
 coat of the intestines abraded in the stools; but Zimmerman declares it to be nothing
 but indurated mucus forced out from an inflamed surface.

The appearance of pus among the stools has by some been considered as a

Proof of a law
but is now well
and joined from
The various things
and which they are
that opinion is
the uniform duty
Others describe
of which they come
land. There have
yet we cannot justify
by relation. Some
but their comparison
The various states
the matter however
sometimes this has
only the effect of
doubt in which it
The various of you
words and acco

Proof of ulceration of the intestines having taken place, but the fallacy of this is now well known, it having been ascertained that pus may be secreted and poured from the surfaces of mucous membranes without any erosion whatever. The earlier physicians speak of substances voided by stool called *Caecumula* and which they considered as pieces of the muscular coat of the intestines. That opinion is probably correct as subsequent experience has ascertained the uniform danger attendant on such casts.

Others describe substances denominatd *Copora Fringuia*, the appearance of which they conceived denoted an affection of the lower part of the alimentary Canal. There have been many vague suppositions as to their formation and structure, yet we cannot possibly conjecture them to be any thing else than the faeces changed by retention. Some have thought the *Caecumula* and *Copora Fringuia* the same but their comparative mortality well establish a distinction.

The various substances evacuated by stool are often thrown up by vomiting. The matter thrown up by vomiting is generally more or less bilious, but what connection this has with the disease we are unable to say, most probably it is merely the effect of the repetition of that process and the particular person and climate in which the disease occurs.

The various symptoms of a disordered state of the alimentary Canal often precede and accompany the disease.

After a short
upon the state of
with the disease
Many other
particularly the
sympathy exist
should on this p
deliberate sta
symptoms.

Recap 1800
as affected, and
later stages.

A painful
has resorted to
but which Mose
produced by the
by the operation of
on the other hand,
circumstances that
when worse it was

Altho sometimes occur, and when they do, are to be considered rather as dependent upon the state of the general system, than as having any particular connexion with the disease.

Many other eruptions are said to be a frequent attendant on this disease particularly the Miliaria, and they are supposed to depend upon the sympathy existing between the surface and alimentary canal. We should on this ground be lead to judge them (when not occurring in a debilitated state of the system or as a consequence of it), as a favorable symptom.

Hiccup sometimes early occurs when the higher parts of the intestines are affected, and this is not alarming, which is always the case in the latter stages.

A painful strangury sometimes attends this disease, which has been ascribed to the sympathy between the rectum and neck of the bladder; but which mostly with more probability we think, supposed always to be produced by the improprie use of opium. This does not preclude the explanation by the operation of any sympathy which may exist between these parts, but on the other hand, the tremendous griping and pain which accompany the evacuations that succeed the liberal use of opium, would call it into action were it ever so slight.

Delirium is
find it (18th)
always precedes
previous secret
Dysentery
times the fever
usually ceases.

There are
death, at some
delirium may de
symptoms. It is
the Dysentery,

Sometimes
stomach, as Dy

Dysentery
by death is fast
When it termin

ten to fourteen

Sometimes it
patients unhappy

Debility is said to predominate in this disease, but we believe we shall find it (taking into view the depression which inflammation of these parts always produce, and excepting the Syphilitic Stasis) rather proportioned to the previous excitement on the lips by evacuations.

Dysentery sometimes terminates as it comes on with Diarrhoea; at other times the frequency of the stools and griping gradually abating, the disease totally ceases.

These are the most usual terminations in health, but when it ends in death, it may be either in a few days by excessive evacuations, or a fatal debility may be produced by a more protracted yet severe state of the symptoms. It also often ends fatally by a bilious purging which succeeds the Dysentery, tho' this is not uniformly a mortal symptom.

Sometimes the disease leaves behind it troublesome affections of the stomach, as Dyspepsia, Flatulence &c. &c.

Dysentery when it ends fatally is often in a few days, but more generally death is produced by the gradual exhaustion of the vital forces.

When it terminates in health, the most usual period of duration is from ten to fourteen or twenty days.

Sometimes it becomes chronic, lasting months, years, and even perhaps the patient's unhappy life; this form is of very difficult cure.

Diagnosis

It is Scarce, and
led down to death
that this object may
of all the symptoms
occasionally in con-
sulted, peculiar

Prognosis

or of the Pleuritic form
must very rarely, and
a dangerous disease
symptoms. When it
may occur.

The symptoms are
and a moderate Pleuritic

The relative degree
the Pleuritic, affords some
doubt.

The terminus and
here and in small

Diagnosis. The only disease with which Dysentery is liable to be confounded, is Diarrhoea, and notwithstanding the several distinctive marks which have been laid down to enable us to distinguish the one from the other, we are humbly of opinion that this object may be better attained, by the consideration of the proportional severity of all the symptoms, which are essential to these two diseases and Typhoid, occasionally in common by them, than by attending to any one which may be supposed peculiar to either.

Prognosis. When there is no fever the disease is mild and when it is slight, or of the Remittent form, it affords a favorable prognosis. When it is in the commonest very severe, as it is followed by a proportional degree of debility, it marks a dangerous disease; the not so much so as when the debility is one of the primary symptoms. When it is of the Typhoid type there is danger at whatever period it may occur.

The softness and moisture of the skin, the appetite still remaining good, and a moderate state of the other symptoms are among our best prognostics.

The relative degree of the tormina and tenesmus and the appearance of the stools, afford our last means of ascertaining the probable termination of the disease.

The tormina and tenesmus being frequent and the faces being voided with much pain and in small quantities are unfavorable. On the contrary altho the faeces

may be some, yet of
with a temporary remedy
When pure blood is
is much danger, the
The presence of gangrene
due to the touch, &c. &c.

Upon the whole it was
to form his prognosis for
he should do it only
The practice of prognosis
is to ascertain the disease
pending active remedies

Morbid Appearances

of active inflammation
Sometimes there are ulcers
intestinal caused even to
be often met with
in find the coats thickened
ten, or putrefaction, or
The sometimes find con

may be severe, yet if there are copious evacuations of the natural feces attended with a temporary remission, we have little to dread.

When feces blood is secreted from the commencement in large quantities, there is much danger, the patient being often carried off in a few days.

The presence of gangrene is denoted by dark, fetid stools, a tense abdomen being due to the touch, &c. &c.

Upon the whole it would be improper and altogether impossible for a practitioner to form his prognosis from the severity or mildness of any one symptom; but he should do it only from the consideration of them all conjointly taken.

The practice of prognosis, as we think above principally, so far as it enables us to ascertain the danger of the disorder, and admonishes us to employ corresponding active remedies for its removal.

Morbid Appearances after Death. We always find marks of active inflammation having existed in the intestines previous to death.

Sometimes there are ulcerations, erosions and sphacelated spots throughout the intestinal canal even to the stomach.

We often meet with portions of a dark colour in a gangrenous state; again we find the coats thickened and soft; the latter on the effects either of inflammation, or putrefaction, or perhaps both.

We sometimes find constrictions of the large intestine, and particularly in

The skin; but
did not submit to
all slight constric-
tions. The
inflammation, and the
the large intestine
lower and farther
primarily affected.
The sometimes fir-
mities, then being
between the sections.
Inflamed, inflaming
in veins, as the stom-
ach, flat, black, is
it as having been for-
mally been detached, the
the intestine after death
was, containing a large
the intestines are often
distended with air.

the Colon; but their non-appearance upon dissection is no proof that they did not subsist before death, for as Hippocrates observes "Mort quodnos soluit," all slight constrictions, all minor inflammations are thus dissipated.

Constrictions show the muscular as well as the internal coat to have been inflamed, and this together with the inflammation being most usually in the large intestines, appear to be the principal differences between this disease and Calculus, in which as we imagine the Villous coat alone is primarily affected.

We sometimes find the inflammation extending to all the coats of the intestines, then being adhesions between their constrictions, gangrenous spots between their external surfaces, abscesses between their coats &c. &c.

Sphacelus, inflammation, or abscess are occasionally found in the neighbouring viscera, as the Stomach, Liver, Spleen, Pancreas, &c. &c.

Small, flat, black tubercles resembling the confluent small Pox, are mentioned as having been found in the internal coats of the intestines; but they have rarely been described, tho' some think this arose from the inattention of dissection. The Mesentery after death is often loaded with fat, tho' sometimes it is excessively wasted, containing abscesses, tubercles &c. &c.

The Intestines are often enormously enlarged, the consequence of previous distention with air.

From the first
been decided,
when it was at a
circumstantial, ha

Causes.

the two kinds of
existing to be co
be seen as how
climate and chie
in how climate
suffered the the
affected by how a
induced by storm
proving that it was
in the bowels and a
from that which we
in kind it's a den
hence from the low
has been the causes of
each of one, and the

From the few cases that have been mentioned where the thoracic viscera have been diseased, we conclude this to have rarely been found the case, and where they are it was at we imagine not as an effect of this disease, but merely an accidental concomitant, having previously existed.

Causes. In detailing the causes of dysentery we shall divide them only into the two heads of proximate and remote; believing the remote predisponent and exciting to be with difficulty separated, and when they do operate they being the same as those of febrile diseases generally. As Dysentery is a disease of warm climates and chiefly occurs in the season superseding the one of the greatest warmth in those climates, we in general terms ascribe its origin to circumstances leading to suppress the the perspiration, acting on the system when it is most liable to be affected by those causes. By Dr. Mosely this disease is supposed always to be produced by obstructed perspiration when it prevails epidemically; at the same time granting that it may be occasionally caused by the operation of accidental stimuli in the bowels, and as his theory does not inculcate a fraction materially different from that which we should deduce from a knowledge of the proximate cause alone, we think its admission unattended by those bad consequences, which hastily adopted theories from the hands of the inexperienced often produce. In our attempts to discover the causes of diseases we can only be directed by the experience of others, the mode of cure, and the consideration of the symptoms as they occur in different stages

of the disease.
In diseases, which
ought to be said
In his disease,
States of the mind
of such an effect
Mental Operations
moral; which
In the first and
by debility, some
the utility of the
time as to their
his influence to
truth, as we can
Imaginations, their
beliefs, they being
not with the view
his disease, and
powerful avail
From his imperfec

of the disease, assisted by a discrimination between those produced by the causes of the disease itself, and those coming from association or sympathy, which may properly
 might be said to arise from the primary disorder action and propagating itself;
 In this disease, we find all practitioners enumerating among its causes those
 states of the weather and seasons of the year, which are most favorable to the production
 of such an effect. This theory also enables us on rational grounds, to account for the
Morbid Operation of the remedies which are so universally recommended for its
 removal; which affords an additional argument in support of its validity.
 In the first and even in more advanced stages, when not counterindicated
 by debility, Emetics are by all strenuously advised and experience confirms
 the utility of the practice without practitioners having them formed any delu-
 sion as to their mode of action. Morely's theory would unhesitatingly ascribe
 their influence to their relaxing effect, and we have here as much proof of its
 truth, as we can have of any other of the same nature. With respect to the
 purgatives, their operation may be thus equally well explained as that of
 emetics, they being both auxiliary to the diaphoretic plan. Diaphoretics do
 not with the view to their sudorific effect, have been long used in the cure of
 this disease, and Morely boldly confides in their use, aided by the most ef-
 fective auxiliaries Elix. and Cathartics.

From the imperfection of our Physiological knowledge, we are unable to

deduct those
which a surface
difficulty which
connection betw
in the case of D
that this is al
that there is som
determining he
as long as we
exist in so ma
with their penne

Mostly with
hasen thereon
is always observ
that he has observ
with those of the
tion of the diffic
in many other
in inability for
or his were one

deduce those facts relative to the causes of a disease from the symptoms, which a superficial view of the subject would lead us to surmise. Another difficulty which opposes itself to us advancement in the development of the connection between the causes and the symptoms of disease, is, as for instance in the case of Dysentery, assuming obstructed perspiration as its cause, the fact that this is also the source of many other diseases; hence we must suppose that there is some conjoint cause, associated with obstructed perspiration determining the diseased action to be bowels in preference to any other parts. As long as we remain ignorant of those conjoint causes, whose influence exists in so many diseases, we must probably so long remain unacquainted with their remote causes.

Mostly with Sydenham supposes Dysentery to be merely the Fever of the season thrown on the bowels, the former still, understanding that the remote cause is always obstructed perspiration. Mostly states in confirmation of his opinion, that he has observed the exacerbations of the disease regularly to correspond with those of the Fever at that time prevalent. Whether this affords any solution of the difficulty above mentioned, we cannot say. In the present case they do in many others, for the reasons already stated, we are compelled to acknowledge our inability from the symptoms of a disease to ascertain its remote causes, yet this were our knowledge perfect, should afford me of our most abundant

[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]

source of information
There are also
dist. information
as the one we have
and account of
is here unnecessary
it is to be proved
We shall not
rather than
have one plan
to anticipate
By the President
discovered action
standing to the
death for need
the President
trial, most ad
there, the court
and the President
appears but where

sources of information on this branch of science.

There are also other causes of Dysentery besides obstructed perspiration, as bad diet, improper articles of food, bad water, &c. &c. whose function is not so extensive, as the one we have been considering. Having in another place spoken of contagion, and avowed our total disbelief in the existence of this as a cause of Dysentery, it is here unnecessary to mention the various means by which authors have thus imagined it to be propagated.

We shall now make some remarks on the Proximate Cause of Dysentery, perhaps rather prematurely, as they may be thought to come in better after we have laid down our plan of treatment; but it is necessary to be ever vigilant, tho' we may have to anticipate some things which will be hereafter mentioned in the proper place. By the proximate cause of a disease, we understand the *ipso motu* the diseased action itself. Now we shall be assisted in our inquiries, by particularly attending to the symptoms, method of cure and the morbid appearances after death, for reasons already given neglecting the remote cause. By Cullen and others, the proximate cause is judged to be a constriction of some part of the alimentary canal, most usually the Colon, and that inflammation supervents upon it. By others, the constriction is said to be a consequence of inflammation. In both cases the primary symptoms of pain and retention of the natural feces may appear, but where there is violent tormina and tenesmus, which often is the

case even in the
irritability which is
found the frequent
little supposition
is an inclined to
which has been for
its truth. By this
perhaps after the
prompt exhibition
ration produced
as the only reason
of the latter, some
of the former. The
have said that, pro-
either one or both
yet we think at any
the arguments de-
prostate cause of
our Pathology of
Shaking, is the Force

case even in the first stages, there can be no doubt as to the existence of an irritability which most unequivocally denoted the presence of inflammation. From the frequent and early appearance of these symptoms, and because the latter supposition best agrees with the remote cause which we have adopted, we are inclined to range ourselves among its advocates. A view of the treatment which has been found most successful tends to corroborate, and goes far to establish its truth. By Cullen's hypothesis, we should be led in the commencement, perhaps after promising it, or even without it, to have recourse to the early and prompt exhibition of Opiated and Antispasmodic: from the theory of inflammation produced by suppressed putrefaction, Emetics, Purgatives, and Cathartics, are the only remedies to which we can trust. Experience has tested the utility of the latter remedies, and the inefficiency and manifestly injurious effects of the former. The appearances on dissection, always show inflammation to have existed previous to death, sometimes constitutions, tho' the existence of either one or both, does not lead us to ascertain which was the primary symptom, yet we think as any inferences drawn from the morbid appearances are equivocal, the arguments deduced from the only other means we possess of discovering the proximate cause of a disease, are abundantly sufficient to establish our theory.

Our Pathology of this disease, is then that; Dysentery when Epidemic strictly speaking, is the Fever of the Colon thrown on the bowels by obstructed putrefaction,

which has been
cut out from
the

And now
to make some
on views of the
which we have
good, we must
path of history
appear to have
other viewed
the labours of
responsibility
Tralmen

passed to the
but suited to a
healthy action
desired, the of
comes when the
lower with the

which produces an inflammation of the intestines, whence originate constipation and from these two diseased actions all the symptoms observed take their rise.

And now having gone thus far, this may not be an inappropriate place to make some apology for the imperfect manner, in which we have elucidated our views of the theory of this disease, and to give some explanation of the standing, which we think them entitled to hold. Unaided by the experience of numerous years, we must when we have deviated from the beaten track, have entered a path of theory directed alone by reason and argument; and when they shall appear to have deserted us, we wish the propositions thus unsupported to be rather viewed in the light of hints and queries, which may serve to assist the labours of the more profound, than as assertions resigned upon our own responsibility.

Treatment. Conformably to that theory we have laid down, we shall proceed to the consideration of that treatment of this disease, which we think best suited to the removal of the remote cause and invigorating the natural healthy action of the parts, after the eradication of the diseased one. In all diseases, the object of our remedies must be either the removal of the remote causes when they are evident; or by the inferences deduced from an acquaintance with them, our endeavours must be directed to the subduing the

passionate
from the im-
passioned
-passion in
from the im-
confess the i-
just and ra-
seconds for
by nature, is
to the restora-
previous offer
-stitute and to
In Dyce
an indication
passionate or
an therefore
to give them
an not directly
rather to be re-
In the case

proximate cause, which amounts to nearly the same. Again, in other cases from the insufficiency of one means, we are compelled to substitute another more powerful diseased action in place of the existing one, and thus we are often successful in the cure of diseases. This latter method is only to be resorted to, when from the impotence of one act, or the imperfection of our knowledge, we have to confess the inadequacy of our remedies to the removal of the disease upon just and rational principles. It is manifestly unnatural, and but an excuse for our ignorance. In other cases where the mode of cure dictated by nature, is dangerous to be pursued, where even when nature succeeds to the restoration of health, but leaves the system so debilitated from the previous effects as exposes the patient to fall its victim; we must here substitute another and a safer one in its place.

In Dysentery we think the remote cause sufficiently manifest to form an indication upon; at the same time paying a proper attention to the proximate cause, so far as we can trace their connection. Our indications are therefore to restore the natural healthy action of the capillaries, and to give them tone. Some of the remedies which we shall use for this purpose, are not directly adapted to the fulfilment of these indications, but are rather to be regarded as essential subsidaries.

In the commencement of the disease, it has long been the practice

to resort to
anti-jerists
Mostly the
hills, which
chiefly for
destroying
by other, of
think that
practised, the
being those of
for as to adve
possession on the
story of all the
necessary but
with a loaded
climate; with
is the historical
useful remedies
In the course
we must resort

to resort to emetics, with the view of cleansing the stomach, and by the anti-peristaltic motion thus excited, to check the action downwards. By Morrell they were used with the intention of evacuating superabundant bile, which from the season and climate was a frequent concomitant; and chiefly for the purpose of causing a revulsion to the surface, and thus at once destroying the disease. By him his Veticolic solution was greatly preferred; by others, Speacuanha and Tartar Emetic were used. We are disposed to think that Emetics were more admissible in the climate in which Dr. Morrell practiced, than they would be found in the United States, from the disease being there often accompanied with bilious diseases. Morrell even goes so far as to advise their repetition, should not the first make a favorable impression on the surface. Dr. Chapman thinks Emetics best suited to Dyspepsia of all the blood affections; but still he conceals that they are not only necessary but indispensable in cases combined with Intermittent Fever, or with a loaded and oppressed stomach, which is often the case in warm climates; with the Professor we here entirely coincide. Our greatest objection to the habitual use of Emetics, is that they interfere too much with the more useful remedies cathartics, of which we shall come presently to speak. In the commencement of the disease, and prior to the use of all other remedies, we must resort to V. and proportion the quantity to the degree of excitement

and local influence
to prepare the system
By all the previous
one opinion, from
society, in consequence
we believe, will be
use of it, that a
remedies in human
state of the system
then. In more or
should be sacrificed
the quantity by the
after after the one
have remedies which
We will meet
in this disease. but
remedies was that
remedies cathartic
of the disease while
practice of giving it

and local inflammation present: This is not only to lessen action, but to prepare the system for the more effectual operation of diaphoretics. By all, the propriety of it in certain cases has been admitted, but in our opinion, practitioners generally have been much too sparing of this remedy, in consequence of their dread of the succeeding debility, which as we believe, will not be so readily produced, by the prompt and active use of it, thus at a blow rooting out the disease, as by tampering with remedies in themselves inefficient or rendered so by the inflammatory state of the system. It is an indispensable auxiliary to the diaphoretic plan. In more advanced stages we should not be deterred from its use, should the excitement or local symptoms require it; only regulating the quantity by the state of the system as well as of the local affection. After the moderate debilitation of blood, the disease will yield to those remedies which before had been ineffectually used.

We will next speak of Cathartics which are of primary importance in this disease. We have already observed that our greatest objection to Emetics was that they too much interfered, with the more necessary remedies cathartics. We will know that we cannot obtain a solution of the disease while the bowels are constipated; and we cannot think the practice of giving Emetics and trusting to Antimonials and Spessminerals

for their own
most active
which is one
cathartic, so
by fertility
in more power
You are in the
to produce a
in expectation
greatly satisfied
when in the con-
tin does not
it will be much
we have advised
For the party
recommended, in
action at some
for delay. The case
anxious to the
has reasons we

for their evacuation will answer so well, as the vigorous use of the most active Cathartics at once. Our practice then is after prescribing ℞. which in most cases will be necessary, to exhibit a large dose of some active cathartic, so as to produce a complete evacuation of the intestines; thus greatly facilitating the operation of the other remedies we may recur to, and which are more particularly adapted to the fulfilment of our indications. Now on the commencement without the previous use of these remedies, attempt to produce a resolution to the surface, we should not only be disappointed in our expectations of cure; but in most cases the disease would most speedily be greatly exacerbated by this imprudent course. No doubt could be raised do occur when in the commencement, diaphoretics would succeed, but as our discrimination does not always enable us to distinguish them with sufficient accuracy, it will be much the safer and more efficient practice to pursue the course we have advised.

For the purpose of emptying the bowels, a variety of articles have been recommended, but we should be induced to pursue the same action as most active as ~~soon~~ promptly fulfilling our object; for here we have no time for delay. The excitement is to be subdued, the bowels emptied, and a resolution to the surface produced, as speedily as circumstances will allow. For these reasons we should always choose Calomel in large doses, assisted by

by letter Feb. 18
salutations of love
all answer, but in
After having ad-
vise to such a
and determining
there is much over
dilemma. By these
but from the con-
and and particu-
undelivered to give
tens of denied to
ing happened. I
I would answer o
It is possible th
Produce some other
Lundamen, might
collected that by
to increase his story, a
just worth attending

by Castor Oil. Perhaps where a repetition of the Purgative is necessary, Oil, combinations of Salt, Senna and Manna which have been much recommended will answer, but not on somewhat prejudiced in favour of Calomel were here.

After having advanced thus far in the treatment of Dysentery, we would next recur to such Diaphoretics as fulfill the double indication of opening the bowels and determining to the surface. For this purpose Spicesnake in nauseating doses is much recommended and appears eminently adapted to attain these objects. By these the preparations of Antimony are used with the same view, but from the concurrent testimony in favour of Spicesnake by many practitioners and particularly from that of our very able Professor Dr. Chapman; we are disposed to give it the decided preference. Dr. Mosely speaks in the highest terms of Sweet Powder as well suited to the evacuation of the bowels, and producing diaphoresis. We are inclined to think that could we procure it pure, it would answer very well.

It is probable that immediately after the operation of the Purgative, Sweet Powder or some other enter diaphoretic, as Antimonial wine combined with Laudanum, might go far towards the removal of the disease. It is to be recollected that by the addition of Laudanum to Diaphoretics, we are enabled to increase the dose, without the inconvenience of exciting nausea. Another fact worth attending to, is that during a prostration, spiritus may be

land in a
Pharm is also
stages of the a
We cannot
Pharm in the
real use of the
some rational
of their medicine
termina which
crises on ad
The flow of
ally have to p
but we have in
quite to avoid
medicines.

Having all
be bound, and
you that the
quant solution
compelled to the

be used in much larger quantities with perfect safety than at other times. Opium is also useful to procure sleep and allay pain in more advanced stages of the disease, when all danger as to its stimulating effects are removed. We cannot give a more luminous and combinatorial view of the use of opium in this disease than by quoting the following from Parry: "The real use of opium is to arrest the hurry of the disease; to procure time to put some rational method of cure into execution; to take off the irritating property of other medicines, and to give them their intended effect and to ease those termina which are sometimes intolerable. Hence the matchless power of opium raises our admiration."

The plan of treatment we have laid down is such, as we shall most usually have to pursue; but in some cases the violence of the symptoms is such, that we have in the beginning to attack the disease with spirits and bleeding to arrest its progress and afford time for the exhibition of other remedies.

Having attained the object of reducing excitement, completely evacuating the bowels, and producing a reaction to the surface, 'tis rational to imagine that the removal of the remote cause would be attended by a consequent solution of the disease; but this is not always the case, and we are compelled to the use of other remedies, which are to be varied according to

The state of
which should
but the knowledge
notion to be placed
of Sarcina, we
of Sarcina and the
the Sarcinae
certain cases of
Chronic S
eating conditions
good habits of
all. My Son, N
good in many
cases by the action
any particular
intended in the
attention by the
Pythius Sta
Pythius, or this fever
materially different

The state of the symptoms, the period of the disease, and other circumstances which should influence our prescriptions. It should however be recollected, that the bowels must be strictly attended to in every stage. When we have reason to suspect the disease to be kept up by habit, or it approaches the form of Dysentery, we may with safety and advantage resort to the cautious exhibition of astringents. Dover's Powder, Kino, Logwood, Cassia, and other astringents and Tonics of a like character are those we should; but in certain cases absorbents might not only be useful but indispensable.

Chronic State. Here we shall derive most benefit from invigorating remedies, a strict attention to diet, and a change of the climate and past habits of the patient. The exercise of gestation is here eminently serviceable. By some, Mercury has been highly extolled; it has certainly done good in many cases, and where we are foiled in our attempts to effect a cure by the ordinary methods we are justifiable in recurring to its use, any particular symptoms not contra-indicating it. It must be gently introduced into the system, the patient being supported during the salivation by the diffusible stimuli.

Typhus State. Here it may be either originally a combination with Typhus, or this fever may supervene upon it as in other diseases. The practice is materially different in these two cases. In the first as in Eriopsatic Typhus we

leave the common
strong and propin-
quity, we must need
help thinking, be-
cause, however, there
by the consideration
and system.
In the second
instance, occur
the strength, which
may be advantage-
ous in itself, but
is inadvisable, be-
cause of opinion and
to think generally of
most money, and
When dysentery
of many, particularly
of the back; and so
declares himself to
one of those cases which

have in the commencement to exhibit Stimuli. After we have by limits made a strong impression on the system, and by cathartics completely evacuated the Primæ Viæ, we must recur to Tonics and Stimulants as in Typhus Febris. You cannot help thinking, but that in this state, there is greater room for purgatives and Diaphoretics, than in Stipathic Typhus. The treatment however will be governed by the consideration of that of the two diseases, and the state of the symptoms and systems.

In the second state when Typhus supervenes upon Dysentery, we must instantly recur to Mercury, gradually insinuating it into the system. The strength must be upheld by the diffusible Stimuli. In many cases it may be advantageous to combine Nitric Acid with it; and when the Mercury is inadmissible, Nitric Acid may be substituted. By some combinations of equal parts of Opium and Nitric Acid have been highly recommended. We are disposed to think favorably of the medicine in those cases where from weakness or other causes Mercury cannot be used.

When Dysentery is combined with Intermittent Fever, it has been the practice of many, particularly Clifton to seize the earliest opportunities for the exhibition of the Bark; and even when there was only slight remissions of the Fever, he declares himself to have derived much benefit from its use, effecting a cure of those cases which had remained intractable under any other mode of

treatment.

Dr. Chas.

directed by the usual
which practice he

Having now our
method of treatment
for us, here to me
the regimen, which

We will begin with
after the excitement
should the pain come

particularly when a
series of blisters are
they are also advantage

may doubt as to the
to the abdomen, be a

relieving pain, and
treatment. In the m.

Next, we may use the
could be in sitting or

treatment.

Dr. Chapman says he is in the habit first of curing the local disease by the usual remedies, and then attacking the fever with the Bath; which practice he has found much more successful than that of Elixophor.

Having now noticed the different forms of Dysentery, and explained the method of treatment which we think best adapted to each, it only now remains for us, here to make some observations on a few particular remedies and the regimen, which we judged to be the most suitable place for them.

We will begin with Blister and other topical applications to the abdomen. After the excitement had been somewhat reduced and the prima via emptied, should the pain continue severe, and then appearing symptoms of inflammation, particularly when in the small intestines or the colon, we would suppose the application of a Blister as near as possible the part affected, would prove of much service. They are also advantageous from their action on the skin. No other cases where there is any doubt, as to the propriety of Blister, fomentations, poultices of warm milk to the abdomen, &c. are grateful applications and would certainly prove useful in allaying pain, and in such cases may be resorted to as a part of our palliative treatment. In the more advanced stages, or even earlier when we want a stimulant effect, we may use Spices, Spirits Turpentine, &c. in Bath &c. Their chief utility would be in aiding in other means of producing perspiration. We are not disposed

to place given
diaphanous for
new from down

Cylinders. A

show the design
and imitation,

vacuumation of the

textures, nor,

materials. Slips

that are perfect

Flannel

Dr. Chapman is

they were by his

find in the sect

ready to essay

chronic states,

effects are to give

to aid respiration

is more adu

cheat possible,

to place great confidence in the Warm Bath, or other external means of exciting diaphoresis farther than as auxiliaries. Used with this view they may in certain cases prove serviceable.

Rhytides. Rhytides are a remedy of much efficacy in every stage of the disease where the symptoms would seem to require them. Their use, is to allay pain and irritation, to loosen the tongue and trachea, and to procure ease in the evacuation of the bowels. They are never to be used with the view of superinducing attraction, nor should they even be composed of saline or any other irritating materials. Infusions of fresh butter, lard, mucilaginous and oily substances that are perfectly bland, or such as we should always prefer.

Flannel Roller. Suggested by Dewar, he previously used by Dr. Chapman in his private practice with the same views as those with which they were by him recommended. Dewar's praises of the remedy are exaggerated in the extreme, in Dysentery and Diarrhoea. He restricts the use of the remedy to every stage of febrile diseases, but Dr. Chapman confines them to the chronic states, when we think them certainly most entitled to credit. Their effects are to give support and tone to the intestines, to protect from cold, and to aid respiration. When applied early in the disease, it should not be so tight as in more advanced stages. In the chronic state it may be applied as often as possible, without impeding respiration. In the first, we expect to

to drain himself
and giving time
should in all cases
think it in some
of Dysentery and
Rugimen.
fascinating sub-
ject as entirely
lines of Quin and
generally used, a
fact of the treatment
-proof of a danger
institutions yet de-
pended enough
acute Dysentery
apostrophe in the case,
under the use of it
It might perhaps
use of wine, Black
and concerning us

to derive benefit from its warmth, in the latter from supporting the abdomen and giving tone to the intestines. It should completely cover the abdomen. We should in all cases be careful not to affect the expiration by it. We are disposed to think it in some cases a useful auxiliary, particularly in the chronic state of Dysentery and Diarrhoea. In our opinion best suited to Diarrhoea.

Regimen. The diet in this disease must be mild and digestible. The farinaceous articles are generally preferred. Decoctions are of great use both as articles of food and of medicine. The juice of the Slippery Elm, solutions of Gum Arabic, Tapioca, Roan Root, Sage, Barley water, &c. are those generally used, and by many they are thought to constitute an indispensable part of the treatment. In more advanced stages when we have to fear the approach of a dangerous debility, Chicken Broth and other articles of more nutritious yet digestible properties are to be used. It may perhaps be properly enough mentioned here that the patient should always in acute Dysentery be confined to a recumbent posture. This often greatly assists in the cure, and neglecting it the disease has often proved obstinate under the use of the usual remedies.

It might perhaps be expected that we should have given our views as to the use of wine, Bark and Mercury of which so much has been said by authors and concerning which there appears to exist such a diversity of opinions.

If Mercury is
nearly clear,
system. On his
own to the
ness of these men
he just will to
is strictly and
According to a
Model of Man
ideal in his party
You have more to
to have been divided
he chooses and
of his kind that
-ed the bounds
-plied our resources
-dition of our species
only he sets first
gentleman to who
flattering remark

Of Mercury we have already said enough. of Wine and Bark, we will merely observe, they are never to be given in the inflammatory state of the system. On this subject we conceive it unnecessary to say more, as a reference to the treatment of Syphilis Venæ, will fully explain our views of the uses of these medicines in this disease. We will only further remark, that the great rule to direct us, and by which being governed, we can rarely err, is strictly and carefully to attend to the state of the system.


According to our promise we would here enter into the consideration of the Modus Operandi of the preceding remedies, had we not so fully given our ideas on this subject in another place, as to leave nothing we could say, unsaid.


We have now brought this paper to a close. Much more might and ought to have been said, to have made it a perfect treatise, to have faithfully exhibited the theories and practice of the ancient and moderns, but from a production of this kind that could neither be expected or required. We have already exceeded the bounds which we assigned ourselves, and perhaps unnecessarily multiplied our remarks. Shall we have succeeded in giving a clear and lucid exposition of our opinions of the theory and treatment of this disease, we shall not only be satisfied first, but will deem such a verdict from the distinguished gentlemen to whom examination this will be referred, as the greatest and most flattering mark of our success, which can possibly be conferred upon us.

We are conscious
and deeply value;
alterations which
improvements is the
more time to this
more perfect by me
that the task has
of our own & criticism
the that we make
from us, but we must
write in his own
deepest criticism of
we confess our close
prejudice of flag, he
The man with diffi-
culty body, whom
we can obtain. There
is, at least it will
to the laws of our culture
qualifications (the an

We are conscious that on some parts we have been obscure, on others perhaps needlessly verbose; in other places our observations are unattended by those explanations which might have accompanied them. Our only excuse for these imperfections is the hurry in which we have written. Could we have devoted more time to this purpose, we flatter ourselves we might have rendered it much more perfect; by making free use of the works of others, but we have always thought that the task thus imposed on us, was for the purpose of detaching the expression of our own sentiments, rather than those of others. Let it not be understood from this that we make any pretensions to originality. We be such vain attempts from us, but we mean that an accurate account of any disease, from the written in his own language, would as we imagine be considered as the surest criterion of his acquaintance with that disease. On the other hand we confess ourselves greatly indebted to the works of several authors for the preceding Essay, but chiefly to the Lectures of Dr. Chapman.

We now with diffidence and hope, submit it to the judgment of that learned body, whom application we shall esteem the most honorable applause we can obtain. Then we stop and whether it be assigned to infamy or oblivion, we trust it will be viewed rather as a hasty production in conformity to the laws of our celebrated University, than as an accurate test of our abilities and qualifications (tho we have no reason to boast) in Medical Science.





58

6

ad